arXiv:1703.06395v1 [math.FA] 19 Mar 2017

SHARP ESTIMATES FOR COMMUTATORS OF BILINEAR OPERATORS ON MORREY TYPE SPACES

DINGHUAI WANG, JIANG ZHOU* AND ZHIDONG TENG

Abstract. Denote by T and I_α the bilinear Calderón-Zygmund operators and bilinear fractional integrals, respectively. In this paper, it is proved that if $b_1, b_2 \in$ CMO (the BMO-closure of $C_c^\infty(\mathbb{R}^n)$), $[\Pi\vec{b}, T]$ and $[\Pi\vec{b}, I_\alpha]$ ($\vec{b} = (b_1, b_2)$) are all the compact operators from $\mathcal{M}^{p_0}_{\vec{P}}$ (the norm of $\mathcal{M}^{p_0}_{\vec{P}}$ is strictly smaller than $2-$fold product of the Morrey norms) to $M^{q_0}_q$ for some suitable indexes p_0, p_1, p_2 and q_0, q. Specially, we also show that if $b_1 = b_2$, then $b_1, b_2 \in$ CMO is necessary for the compactness of $[\Pi\vec{b}, I_\alpha]$ on Morrey space.

1. Introduction

The aim of the present paper is first: to obtain the boundedness and compactness of iterated commutators of bilinear operators acting on multi-Morrey spaces (a multi-Morrey norm is strictly smaller than $m-$fold product of the Morrey norms); and second: to characterize the compactness of the iterated commutators of bilinear fractional integral operators on Morrey spaces.

A well known result of Coifman, Rochberg and Weiss [19] states that the commutator

$$[b, T](f) = bT(f) - T(bf)$$

is bounded on some L^p, $1 < p < \infty$, if and only if $b \in$ BMO, where T be the classical Calderón-Zygmund operator. In 1978, Uchiyama [37] refined the boundednss results on the commutator to compactness. This is a achieved by requiring the commutator with symbol to be in CMO, which is the closure in BMO of the space of C^∞ functions with compact support. In recent years, the compactness of commutators has been extensively studied already, Wang [39] showed that the compactness of commutator of fractional integral operator and Ding et al. [9], [10], [11], [12], [13] [14] also considered the compactness of commutators for some operators, such as the Riesz potential, singular integral,

2010 Mathematics Subject Classification. Primary 42B20, 47B07; Secondary: 42B99,47G99.

Key words and phrases. Bilinear Calderón-Zygmund operator; Bilinear fractional integral operator; Characterization; Compactness; Commutator

The research was supported by National Natural Science Foundation of China (Grant No.11661075 and No. 11271312).

* Corresponding author, zhoujiang@xju.edu.cn.

Marcinkiewicz integral in Morrey spaces. The interest in the compactness of commutators in complex analysis is from the connection between the commutators and the Hankel-type operators. In fact, the authors of [28] and [29] have applied commutator theory to give a compactness characterization of Hankel operators on holomorphic Hardy spaces $H^2(D)$, where D is a bounded, strictly pseudoconvex domain in $\mathbb{C}^n$. It is perhaps for this important reason that the compactness of commutators attracted ones attention among researchers in PDEs.

Recently, many authors are interested in the multilinear setting, see [5], [21], [22], [23] and [31]. The multilinear Calderón-Zygmund theory originated in the works of Coifman and Meyer in the 70s, see e.g.[17], [18]. Later on the topic was retaken by several authors; including Christ and Journé [15], Kenig and Stein [27] and Grafakos and Torres [23]. The boundedness results for commutators with symbols in BMO started to receive attention only a few years ago, see [30], [32], [33] or [36]. Compactness results in the multilinear setting have just began to be studied. Bényi et al. [3], [4] and [6] showed that symbols in CMO again produce compact commutators. Ding and Mei [20] consider the compactness of linear commutator of bilinear operators from product of Morrey spaces to Morrey spaces. In this paper, some sharp estimates for compactness of commutators of bilinear operators will be given; that is, it is proved that bilinear operators are all compact operators from multi-Morrey spaces(precise definition is given in the next secion) to Morrey spaces.

Another subject of this paper is to consider the characterization of compactness of the iterated commutator of bilinear fractional integral operators. For linear fractional integrals, the characterization of boundedness of the commutator was obtained by Chanillo [8], while the one for compactness is credited in [12] and [39]. In the bilinear setting, in 2015, Chaffee and Torres [7] characterized the compactness of the linear commutators of bilinear fractional integral operators acting on product of Lebesgue spaces. In [38], we obtain the characterization of compactness of iterated commutators of bilinear fractional integral operators acting on product of Lebesgue spaces. In this paper, we will show that CMO in fact characterizes compactness on Morrey spaces.

2. Preliminaries and Main results

2.1. **Bilinear Calderón-Zygmund operator and its commutator.** Recall that bilinear singular integral operator T is a bounded operator which satisfies

$$\|T(f_1, f_2)\|_{L^p} \leq C\|f_1\|_{L^{p_1}}\|f_2\|_{L^{p_2}},$$

for some $1 < p_1, p_2 < \infty$ with $1/p = 1/p_1 + 1/p_2$ and the function K, defined off the diagonal $y_0 = y_1 = y_2$ in $(\mathbb{R}^n)^{2+1}$, satisfies the conditions as follow:

(1) The function K satisfies the size condition.

$$|K(x, y_1, y_2)| \leq \frac{C}{\big(|x - y_1| + |x - y_2| + |y_1 - y_2|\big)^{2n}};$$

(2) The function K satisfies the regularity condition. For some $\gamma > 0$, if $|x - x'| \leq \frac{1}{2}\max\{|x - y_1|, |x - y_2|, |x - y_2|\}$

$$|K(x, y_1, y_2) - K(x', y_1, y_2)| \leq \frac{C|x - x'|^{\gamma}}{\big(|x - y_1| + |x - y_2| + |y_1 - y_2|\big)^{2n+\gamma}};$$

if $|y_1 - y_1'| \leq \frac{1}{2}\max\{|x - y_1|, |x - y_2|, |y_1 - y_2|\}$

$$|K(x, y_1, y_2) - K(x', y_1, y_2)| \leq \frac{C|y_1 - y_1'|^{\gamma}}{\big(|x - y_1| + |x - y_2| + |y_1 - y_2|\big)^{2n+\gamma}};$$

if $|y_2 - y_2'| \leq \frac{1}{2}\max\{|x - y_1|, |x - y_2|, |y_1 - y_2|\}$

$$|K(x, y_1, y_2) - K(x', y_1, y_2)| \leq \frac{C|y_2 - y_2'|^{\gamma}}{\big(|x - y_1| + |x - y_2| + |y_1 - y_2|\big)^{2n+\gamma}};$$

(3) If $x \notin \mathrm{supp} f_1 \bigcap \mathrm{supp} f_2$, then

$$T(f_1, f_2)(x) = \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} K(x, y_1, y_2) f_1(y_1) f_2(y_2) dy_1 dy_2.$$

It was shown that in [23] that if $\frac{1}{r_1} + \frac{1}{r_2} = \frac{1}{r}$, then an bilinear Calderón-Zygmund operator satisies

$$T : L^{r_1} \times L^{r_2} \to L^{r}$$

when $1 < r_1, r_2 < \infty$ and

$$T : L^{r_1} \times L^{r_2} \to L^{r,\infty}$$

when $1 \leq r_1, r_2 < \infty$. In particular

$$T : L^{1} \times L^{1} \to L^{1/2,\infty}.$$

In 2003, Pérez and Torres in [33] defined the commutator $[\Pi\vec{b}, T]$ as follows

$$\begin{aligned}&[\Pi\vec{b}, T](f_1, f_2)(x) := [b_2, [b_1, T]_1]_2(f_1, f_2)(x)\\ &:= \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} (b_1(x) - b_1(y_1))(b_2(x) - b_2(y_2)) K(x, y_1, y_2) f_1(y_1) f_2(y_2) dy_1 dy_2.\end{aligned}$$

They also proved that if $b_1, b_2 \in \mathrm{BMO}$, then

$$[\Pi\vec{b}, T] : L^{r_1} \times L^{r_2} \to L^{r}$$

when $1 < r_1, r_2 < \infty$ with $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$.

The maximal operator T_* of bilinear Calderon-Zygmund operator T is defined by

$$T_*(f_1, f_2)(x) = \sup_{\delta>0} \left| \int_{|x-y_1|+|x-y_2|>\delta} K(x, y_1, y_2) f_1(y_1) f_2(y_2) dy_1 dy_2 \right|.$$

In 2002, Grafakos and Torres in [23] proved that

$$T_* : L^{r_1} \times L^{r_2} \to L^r$$

when $1 < r_1, r_2 < \infty$ with $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$.

2.2. **Bilinear fractional integral operator and its commutator.** It is well known that the fractional integral $\mathcal{I}_\alpha$ of order $\alpha (0 < \alpha < n)$ plays an important role in harmonic analysis, PDE and potential theory (see [35]). Recall that $\mathcal{I}_\alpha$ is defined by

$$\mathcal{I}_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha}} dy.$$

For the bilinear case, the bilinear fractional integral operator I_α, $0 < \alpha < 2n$, is defined by

$$I_\alpha(f_1, f_2)(x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{f_1(y_1) f_2(y_2)}{\big(|x-y_1| + |x-y_2|\big)^{2n-\alpha}} dy_1 dy_2.$$

In this paper, we will consider the following equivalent operator

$$I_\alpha(f_1, f_2)(x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{f_1(y_1) f_2(y_2)}{\big(|x-y_1|^2 + |x-y_2|^2\big)^{n-\alpha/2}} dy_1 dy_2.$$

Its iterated commutator with $\vec{b} = (b_1, b_2)$ is given by

$$\begin{aligned}
&[\Pi\vec{b}, I_\alpha](f_1, f_2)(x) := [b_2, [b_1, I_\alpha]_1]_2(f_1, f_2)(x) \\
&= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{(b_1(x) - b_1(y_1))(b_2(x) - b_2(y_2)) f_1(y_1) f_2(y_2)}{\big(|x-y_1|^2 + |x-y_2|^2\big)^{n-\alpha/2}} dy_1 dy_2.
\end{aligned}$$

2.3. **Morrey type spaces.** The Morrey space was defined by Morrey [31] in 1938, which is connected to certain problems in elliptic PDE. Later, the Morrey space was found to have many important applications to the Navier-Stokes equations [26], the Schrödinger equations [34] and the potential analysis [1] and [2].

Let $1 \le p \le p_0 < \infty$. The Morrey space $M_p^{p_0}$ is defined by the norm

$$\|f\|_{M_p^{p_0}} := \sup_Q |Q|^{1/p_0} \Big(\frac{1}{|Q|} \int_Q |f(x)|^p dx \Big)^{1/p} < \infty.$$

In 2012, Iida et al. [25] introduced the multi-Morrey norm as follow

$$\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}:=\sup_Q|Q|^{1/p_0}\prod_{i=1}^2\Big(\frac{1}{|Q|}\int_Q|f_i(x)|^{p_i}dx\Big)^{1/p_i}<\infty.$$

They showed that Multi-Morrey norm is strictly smaller than 2−fold product of the Morrey norms. They also proved that

$$T:\mathcal{M}^{p_0}_{\vec{P}}\to M^{p_0}_p \text{ and } I_\alpha:\mathcal{M}^{p_0}_{\vec{P}}\to M^{q_0}_q$$

for some suitable indexes p_0,p_1,p_2 and q_0,q. In this paper, we will consider the boundedness and compactness of the commutators $[\Pi\vec{b},T]$ and $[\Pi\vec{b},I_\alpha]$.

2.4. **Main results.** Now we return to our main results.

Theorem 2.1. *Let $\vec{P}=(p_1,p_2)$, $1<p_1,p_2<\infty$, $0<p\le p_0<\infty$ with $\frac{1}{p}=\frac{1}{p_1}+\frac{1}{p_2}$. Suppose that T be a bilinear Calderon-Zygmund operator and $\vec{b}=(b_1,b_2)$ with $b_1,b_2\in$ CMO, then $[\Pi\vec{b},T]$ is a compact operator from $\mathcal{M}^{p_0}_{\vec{P}}$ to $M^{p_0}_q$.*

Theorem 2.2. *Let $0<\alpha<2n,\vec{P}=(p_1,p_2),1<p_1,p_2<\infty,0<p\le p_0<\infty,0<q\le q_0<\infty$ such that $\frac{1}{p}=\frac{1}{p_1}+\frac{1}{p_2}$, $\frac{1}{q_0}=\frac{1}{p_0}-\frac{\alpha}{n}$ and $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$. For the local integral functions b_1,b_2 and $\vec{b}=(b_1,b_2)$, we have*

(1) *if $b_1,b_2\in$ CMO, then $[\Pi\vec{b},I_\alpha]$ is a compact operator from $\mathcal{M}^{p_0}_{\vec{P}}$ to $M^{q_0}_q$.*
(2) *if $b_1=b_2$ and $[\Pi\vec{b},I_\alpha]$ is a compact operator from $\mathcal{M}^{p_0}_{\vec{P}}$ to $M^{q_0}_q$, then $b_1,b_2\in$ CMO.*

3. Main lemmas

To prove Theorem 2.1 and Theorem 2.2, we need the following results.

Lemma 3.1. *Let $1\le p<p_0<\infty$, $1<p_1,p_2<\infty$ with $\frac{1}{p}=\frac{1}{p_1}+\frac{1}{p_2}$. Suppose that T be a bilinear Calderon-Zygmund operator and $\vec{b}=(b_1,b_2)$ with $b_1,b_2\in$ BMO, then*

$$\|[\Pi\vec{b},T]\|_{M^{p_0}_p}\lesssim\|b_1\|_{\rm BMO}\|b_2\|_{\rm BMO}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

Proof. Without loss of generality, we may assume that $\|b_1\|_{\rm BMO}=\|b_2\|_{\rm BMO}=1$. Fixing $Q:=Q(x_0,r)$, we split f_i into $f_i^0+f_i^\infty$ with $f_i^0=f_i\chi_{2Q}$ and $f_i^\infty=f_i\chi_{(2Q)^c}$, $i=1,2$. Then we need to verify the following inequalities:

$$\mathrm{I}_1:=|Q|^{1/p_0-1/p}\bigg(\int_Q\Big|[\Pi\vec{b},T](f_1^0,f_2^0)(x)\Big|^p dx\bigg)^{1/p}\lesssim\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}};\tag{3.1}$$

$$\mathrm{I}_2:=|Q|^{1/p_0-1/p}\bigg(\int_Q\Big|[\Pi\vec{b},T](f_1^0,f_2^\infty)(x)\Big|^p dx\bigg)^{1/p}\lesssim\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}};\tag{3.2}$$

$$\mathrm{I}_3 := |Q|^{1/p_0-1/p}\Big(\int_Q \Big|[\Pi\vec{b},T](f_1^\infty,f_2^0)(x)\Big|^p dx\Big)^{1/p} \lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}; \tag{3.3}$$

$$\mathrm{I}_4 := |Q|^{1/p_0-1/p}\Big(\int_Q \Big|[\Pi\vec{b},T](f_1^\infty,f_2^\infty)(x)\Big|^p dx\Big)^{1/p} \lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}; \tag{3.4}$$

We analyze each term separately. First, we give the proof of Eq. (3.1). The boundedness of $[\Pi\vec{b},T]$ from $L^{p_1}\times L^{p_2}$ to L^p gives

$$\begin{aligned}
\mathrm{I}_1 &\lesssim |Q|^{1/p_0-1/p}\prod_{i=1}^2\Big(\int_{\mathbb{R}^n}|f_i^0(x)|^{p_i}dx\Big)^{1/p_i}\\
&= |Q|^{1/p_0-1/p}\prod_{i=1}^2\Big(\int_{2Q}|f_i(x)|^{p_i}dx\Big)^{1/p_i}\\
&\lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.
\end{aligned}$$

To estimate I_2, the operator $[\Pi\vec{b},T]$ can be devided into the following parts:

$$\begin{aligned}
&[\Pi\vec{b},T](f_1^0,f_2^\infty)(x)\\
&= (b_1(x)-b_{1,2Q})(b_2(x)-b_{2,2Q})T(f_1^0,f_2^\infty)(x) + (b_2(x)-b_{2,2Q})T\big((b_{1,2Q}-b_1)f_1^0,f_2^\infty\big)(x)\\
&\quad +(b_1(x)-b_{1,2Q})T\big(f_1^0,(b_{2,2Q}-b_2)f_2^\infty\big)(x) + T\big((b_{1,2Q}-b_1)f_1^0,(b_{2,2Q}-b_2)f_2^\infty\big)(x)\\
&=: \mathrm{I}_{21}+\mathrm{I}_{22}+\mathrm{I}_{23}+\mathrm{I}_{24},
\end{aligned}$$

where $b_{i,2Q}=\frac{1}{|2Q|}\int_{2Q}b_i(x)dx$ for $i=1,2$.

Now, we give the estimates for $\mathrm{I}_{21},\mathrm{I}_{22},\mathrm{I}_{23},\mathrm{I}_{24}$, respectively. By the definition of T, we have

$$\begin{aligned}
|T(f_1^0,f_1^\infty)(x)| &\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n}}dy_2dy_1\\
&\lesssim \int_{2Q}|f_1(y_1)|dy_1\sum_{k=1}^\infty\frac{1}{|2^kQ|^2}\int_{2^{k+1}Q\setminus 2^kQ}|f_2(y_2)|dy_2\\
&\lesssim \sum_{k=1}^\infty|2^{k+1}Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\
&\lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.
\end{aligned}$$

Similar estimate gives

$$\begin{aligned}
&|T((b_1-b_{1,2Q})f_1^0,f_2^\infty)(x)|\\
&\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q}\frac{|b_1(y_1)-b_{1,2Q}||f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n}}dy_2dy_1
\end{aligned}$$

$$\lesssim \int_{2Q} |b_1(y_1) - b_{1,2Q}||f_1(y_1)|dy_1 \sum_{k=1}^{\infty} \frac{1}{|2^k Q|^2} \int_{2^{k+1}Q\setminus 2^k Q} |f_2(y_2)|dy_2$$
$$\lesssim \left(\frac{1}{|2Q|}\int_{2Q} |b_1(y_1) - b_{1,2Q}|^{p_1'} dy_1\right)^{1/p_1'} \sum_{k=1}^{\infty} |2^{k+1}Q|^{-1/p_0} \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}$$
$$\lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

From the fact that for $b \in \mathrm{BMO}$,

$$|b_{2^k Q} - b_Q| \lesssim k\|b\|_{\mathrm{BMO}},$$

which implies that

$$|T(f_1^0, (b_2 - b_{2,2Q})f_2^\infty)(x)|$$
$$\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q} \frac{|b_2(y_2) - b_{2,2Q}||f_1(y_1)||f_2(y_2)|}{(|x-y_1| + |x-y_2|)^{2n}} dy_2 dy_1$$
$$\lesssim \int_{2Q} |f_1(y_1)|dy_1 \sum_{k=1}^{\infty} \frac{1}{|2^k Q|^2} \int_{2^{k+1}Q\setminus 2^k Q} |b_2(y_2) - b_{2,2Q}||f_2(y_2)|dy_2$$
$$\lesssim \int_{2Q} |f_1(y_1)|dy_1 \sum_{k=1}^{\infty} \frac{1}{|2^k Q|^2} \int_{2^{k+1}Q} \big(|b_2(y_2) - b_{2,2^{k+1}Q}| + |b_{2,2^{k+1}Q} - b_{2,2Q}|\big)|f_2(y_2)|dy_2$$
$$\lesssim \sum_{k=1}^{\infty} k|2^{k+1}Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}$$
$$\lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

Finally, it remains to prove

$$|T((b_1 - b_{1,2Q})f_1^0, (b_2 - b_{2,2Q})f_2^\infty)(x)| \lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

Note that

$$|T((b_1 - b_{1,2Q})f_1^0, (b_2 - b_{2,2Q})f_2^\infty)(x)|$$
$$\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q} \frac{|b_1(y_1) - b_{1,2Q}||b_2(y_2) - b_{2,2Q}||f_1(y_1)||f_2(y_2)|}{(|x-y_1| + |x-y_2|)^{2n}} dy_2 dy_1$$
$$\lesssim \int_{2Q} |b_1(y_1) - b_{1,2Q}||f_1(y_1)|dy_1 \sum_{k=1}^{\infty} \frac{1}{|2^k Q|^2} \int_{2^{k+1}Q\setminus 2^k Q} |b_2(y_2) - b_{2,2Q}||f_2(y_2)|dy_2$$
$$\lesssim \sum_{k=1}^{\infty} k|2^{k+1}Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}$$
$$\lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

Thus, Minkowski's inequality and Hölder's inequality give that

$$\begin{aligned} \mathrm{I}_2 &= |Q|^{1/p_0-1/p}\bigg(\int_Q |\mathrm{I}_{21}+\mathrm{I}_{22}+\mathrm{I}_{23}+\mathrm{I}_{24}|^p\bigg)^{1/p} \\ &\lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}. \end{aligned}$$

We complete the proof of (3.2).

With the same idea of estimate for I_2, we can obtain the similar result for I_3.

To prove (3.4), we need only to show the following four inequalities.

$$|T(f_1^\infty,f_2^\infty)(x)| \lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}} \tag{3.5}$$

$$|T((b_1-b_{1,2Q})f_1^\infty,f_2^\infty)(x)| \lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}} \tag{3.6}$$

$$|T(f_1^\infty,(b_2-b_{2,2Q})f_2^\infty)(x)| \lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}} \tag{3.7}$$

$$|T((b_1-b_{1,2Q})f_1^\infty,(b_2-b_{2,2Q})f_2^\infty)(x)| \lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}} \tag{3.8}$$

Because (3.6), (3.7), (3.8) are completely analogous to (3.5), with a small difference, we only estimate (3.5). Set $\Omega_0 := \{(y_1,y_2) : |x-y_1|+|x-y_2| \le 2r\}$ and $\Omega_k := \{(y_1,y_2) : 2^k r < |x-y_1|+|x-y_2| \le 2^{k+1}r\}$, $k=1,2,\cdots$. Then

$$\begin{aligned} |T(f_1^\infty,f_2^\infty)(x)| &\lesssim \int_{\mathbb{R}^n\setminus 2Q}\int_{\mathbb{R}^n\setminus 2Q}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n}}dy_1dy_2 \\ &\lesssim \iint_{\mathbb{R}^{2n}\setminus\Omega_0}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n}}dy_1dy_2 \\ &\lesssim \sum_{k=1}^\infty\iint_{\Omega_k}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n}}dy_1dy_2 \\ &\lesssim \sum_{k=1}^\infty\frac{1}{(2^kr)^{2n}}\prod_{i=1}^2\int_{2^{k+1}B}|f_i(y_i)|dy_i \\ &\lesssim \sum_{k=1}^\infty|2^kQ|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}} \\ &\lesssim |Q|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}, \end{aligned}$$

where $B := B(x,r)$.

Combining the estimates above, we have

$$\|[\Pi\vec{b},T](f_1,f_2)\|_{M^{p_0}_p} \lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

We complete the proof of Lemma 3.1. □

Now, we give the $M_p^{p_0}-$boundedness for a general bi-sublinear operator which satisfies some control conditions.

Lemma 3.2. *Let S is a bi-sublinear operator satisfies*

$$|S(f_1,f_2)(x)| \lesssim \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|f_1(y_1)f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n}}dy_1dy_2,$$

and for $1\le p<\infty$, $1<p_1,p_2<\infty$ with $\frac{1}{p}=\frac{1}{p_1}+\frac{1}{p_2}$, S is bounded from $L^{p_1}\times L^{p_2}$ to L^p. Then for $1\le p<p_0<\infty$ and $\vec{P}=(p_1,p_2)$, S is bounded from $\mathcal{M}_{\vec{P}}^{p_0}$ to $M_p^{p_0}$; that is

$$\|S(f_1,f_2)\|_{M_p^{p_0}} \lesssim \|(f_1,f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}}.$$

Proof. Fixing $Q:=Q(x_0,r)$ and we write

$$f_i = f_i\chi_{2Q} + \sum_{k=1}^{\infty} f_i\chi_{2^{k+1}Q\setminus 2^kQ} =: \sum_{k=0}^{\infty} f_i^k.$$

Then the L^p-boundedness of S yields

$$\begin{aligned}|Q|^{1/p_0-1/p}\bigg(\int_Q |S(f_1^0,f_2^0)(x)|^p dx\bigg)^{1/p} &\lesssim |Q|^{1/p_0-1/p}\|f_1^0\|_{L^{p_1}}\|f_2^0\|_{L^{p_2}}\\ &\lesssim \|(f_1,f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}}.\end{aligned}$$

To complete the proof of Lemma 3.2, it remains to show the following four inequalities.

$$\sum_{k=1}^{\infty}\bigg(\int_Q |S(f_1^0,f_2^k)(x)|^p dx\bigg)^{1/p} \lesssim |Q|^{1/p-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}}; \tag{3.9}$$

$$\sum_{j=1}^{\infty}\bigg(\int_Q |S(f_1^j,f_2^0)(x)|^p dx\bigg)^{1/p} \lesssim |Q|^{1/p-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}}; \tag{3.10}$$

$$\sum_{j=1}^{\infty}\sum_{k=1}^{j}\bigg(\int_Q |S(f_1^j,f_2^k)(x)|^p dx\bigg)^{1/p} \lesssim |Q|^{1/p-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}}; \tag{3.11}$$

$$\sum_{j=1}^{\infty}\sum_{k=j}^{\infty}\bigg(\int_Q |S(f_1^j,f_2^k)(x)|^p dx\bigg)^{1/p} \lesssim |Q|^{1/p-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}}; \tag{3.12}$$

By the symmetry, we need only to prove (3.9) and (3.11). First, we give the proof of (3.9). Note that

$$|S(f_1^0,f_2^k)(x)| \lesssim \int_{2Q}\int_{2^{k+1}Q\setminus 2^kQ} \frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n}}dy_1dy_2$$

$$\begin{aligned} &\lesssim |2^kQ|^{-2}\int_{2^{k+1}Q}\int_{2^{k+1}Q}|f_1(y_1)||f_2(y_2)|dy_1dy_2\\ &\lesssim |2^kQ|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}, \end{aligned}$$

which implies that

$$\begin{aligned} \sum_{k=1}^{\infty}\bigg(\int_Q|S(f_1^0,f_2^k)(x)|^pdx\bigg)^{1/p} &\lesssim |Q|^{1/p}\sum_{k=1}^{\infty}|2^kQ|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\ &\lesssim |Q|^{1/p-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}. \end{aligned}$$

For $j\geq k$, we also have

$$S(f_1^j,f_2^k)(x) \lesssim |2^jQ|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

Thus,

$$\begin{aligned} \sum_{j=1}^{\infty}\sum_{k=1}^{j}\bigg(\int_Q|S(f_1^j,f_2^k)(x)|^pdx\bigg)^{1/p} &\lesssim |Q|^{1/p}\sum_{j=1}^{\infty}j|2^jQ|^{-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\ &\lesssim |Q|^{1/p-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}. \end{aligned}$$

Thus, we complete the proof of the Lemma 3.2. □

Since bilinear maximal Calderon-Zygmund operator T_* satisfies the condition as in Lemma 3.2, we get immediately the sharp bounds for T_* on Morrey spaces.

Corollary 3.1. *Let $\vec{P}=(p_1,p_2), 1\leq p<p_0<\infty$, $1<p_1,p_2<\infty$ with $\frac{1}{p}=\frac{1}{p_1}+\frac{1}{p_2}$. Suppose that T_* be a bilinear maximal Calderon-Zygmund operator, then*

$$\|T_*(f_1,f_2)\|_{M^{p_0}_p}\lesssim\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

Lemma 3.3. *Under the hypotheses of Theorem 2.2. For the local integral functions b_1,b_2 and $\vec{b}=(b_1,b_2)$, we have*

(1) *if $b_1,b_2\in$ BMO, then $[\Pi\vec{b},I_\alpha]$ is a bounded operator from $\mathcal{M}^{p_0}_{\vec{P}}$ to $M^{q_0}_q$.*
(2) *if $b_1=b_2$ and $[\Pi\vec{b},I_\alpha]$ is a bounded operator from $\mathcal{M}^{p_0}_{\vec{P}}$ to $M^{q_0}_q$, then $b_1,b_2\in$ BMO.*

Proof. Assume that $b_1,b_2\in$ BMO. For any cube Q, we split f_i into $f_i^0+f_i^\infty$ with $f_i^0=f_i\chi_{2Q}$ and $f_i^\infty=f_i\chi_{(2Q)^c}$, $i=1,2$. Then we need to verify the following inequalities:

$$\mathrm{J}_1:=|Q|^{1/q_0-1/q}\bigg(\int_Q\Big|[\Pi\vec{b},I_\alpha](f_1^0,f_2^0)(x)\Big|^qdx\bigg)^{1/q}\lesssim\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}; \tag{3.13}$$

$$\mathrm{J}_2:=|Q|^{1/q_0-1/q}\bigg(\int_Q\Big|[\Pi\vec{b},I_\alpha](f_1^0,f_2^\infty)(x)\Big|^{q_0}dx\bigg)^{1/q_0}\lesssim\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}; \tag{3.14}$$

$$\mathrm{J}_3 := |Q|^{1/q_0-1/q}\bigg(\int_Q \Big|[\Pi\vec{b}, I_\alpha](f_1^\infty, f_2^0)(x)\Big|^{q_0}dx\bigg)^{1/q_0} \lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}; \tag{3.15}$$

$$\mathrm{J}_4 := |Q|^{1/q_0-1/q}\bigg(\int_Q \Big|[\Pi\vec{b}, I_\alpha](f_1^\infty, f_2^\infty)(x)\Big|^{q_0}dx\bigg)^{1/q_0} \lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}. \tag{3.16}$$

By the boundedness of $[\Pi\vec{b}, I_\alpha]$ from $L^{p_1}\times L^{p_2}$ to L^q, we have

$$\begin{aligned}
\mathrm{J}_1 &\lesssim |Q|^{1/q_0-1/q}\prod_{i=1}^2\bigg(\int_{\mathbb{R}^n}|f_i^0(x)|^{p_i}dx\bigg)^{1/p_i}\\
&\lesssim |Q|^{1/p_0-1/p}\prod_{i=1}^2\bigg(\int_{2Q}|f_i(x)|^{p_i}dx\bigg)^{1/p_i}\\
&\lesssim \|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.
\end{aligned}$$

The terms $\mathrm{J}_2, \mathrm{J}_3, \mathrm{J}_4$ are estimates, with slight changes, using the same tools as in the proof for $[\Pi\vec{b}, T]$. For example, if we consider the J_2 term, we first give the estimates for some operators. First,

$$\begin{aligned}
|I_\alpha(f_1^0, f_1^\infty)(x)| &\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|^2+|x-y_2|^2)^{n-\alpha/2}}dy_2dy_1\\
&\lesssim \int_{2Q}|f_1(y_1)|dy_1\sum_{k=1}^\infty\frac{1}{|2^kQ|^{2-\alpha/n}}\int_{2^{k+1}Q\setminus 2^kQ}|f_2(y_2)|dy_2\\
&\lesssim \sum_{k=1}^\infty|2^{k+1}Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\
&\lesssim |Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.
\end{aligned}$$

Second,

$$\begin{aligned}
&|I_\alpha((b_1-b_{1,2Q})f_1^0, f_2^\infty)(x)|\\
&\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q}\frac{|b_1(y_1)-b_{1,2Q}||f_1(y_1)||f_2(y_2)|}{(|x-y_1|^2+|x-y_2|^2)^{n-\alpha/2}}dy_2dy_1\\
&\lesssim \int_{2Q}|b_1(y_1)-b_{1,2Q}||f_1(y_1)|dy_1\sum_{k=1}^\infty\frac{1}{|2^kQ|^{2-\alpha/n}}\int_{2^{k+1}Q\setminus 2^kQ}|f_2(y_2)|dy_2\\
&\lesssim \bigg(\frac{1}{|2Q|}\int_{2Q}|b_1(y_1)-b_{1,2Q}|^{p_1'}dy_1\bigg)^{1/p_1'}\sum_{k=1}^\infty|2^{k+1}Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\
&\lesssim \|b_1\|_{\mathrm{BMO}}|Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.
\end{aligned}$$

Third,

$$
\begin{aligned}
&|I_\alpha(f_1^0,(b_2-b_{2,2Q})f_2^\infty)(x)|\\
&\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q}\frac{|b_2(y_2)-b_{2,2Q}||f_1(y_1)||f_2(y_2)|}{(|x-y_1|^2+|x-y_2|^2)^{n-\alpha/2}}dy_2dy_1\\
&\lesssim \int_{2Q}|f_1(y_1)|dy_1\sum_{k=1}^{\infty}\frac{1}{|2^kQ|^{2-\alpha/n}}\int_{2^{k+1}Q\setminus 2^kQ}|b_2(y_2)-b_{2,2Q}||f_2(y_2)|dy_2\\
&\lesssim \sum_{k=1}^{\infty}|2^{k+1}Q|^{\alpha/n-1/p_0}\left(\frac{1}{|2^{k+1}Q|}\int_{2^{k+1}Q}|b_2(y_2)-b_{2,2Q}|^{p_2'}dy_2\right)^{1/p_2'}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\
&\lesssim \|b_2\|_{\mathrm{BMO}}\sum_{k=1}^{\infty}k|2^{k+1}Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\
&\lesssim \|b_2\|_{\mathrm{BMO}}|Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.
\end{aligned}
$$

Finally,

$$
\begin{aligned}
&|I_\alpha((b_1-b_{1,2Q})f_1^0,(b_2-b_{2,2Q})f_2^\infty)(x)|\\
&\lesssim \int_{2Q}\int_{\mathbb{R}^n\setminus 2Q}\frac{|b_1(y_1)-b_{1,2Q}||f_1(y_1)||b_2(y_2)-b_{2,2Q}||f_1(y_1)||f_2(y_2)|}{(|x-y_1|^2+|x-y_2|^2)^{n-\alpha/2}}dy_2dy_1\\
&\lesssim \int_{2Q}|b(y_1)-b_{2Q}||f_1(y_1)|dy_1\sum_{k=1}^{\infty}\frac{1}{|2^kQ|^{2-\alpha/n}}\int_{2^{k+1}Q\setminus 2^kQ}|b_2(y_2)-b_{2,2Q}||f_2(y_2)|dy_2\\
&\lesssim \left(\frac{1}{|2Q|}\int_{2Q}|b(y_1)-b_{2Q}|^{p_1'}dy_1\right)^{1/p_1'}\|b_2\|_{\mathrm{BMO}}\sum_{k=1}^{\infty}k|2^{k+1}Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\\
&\lesssim \|b_1\|_{\mathrm{BMO}}\|b_2\|_{\mathrm{BMO}}|Q|^{\alpha/n-1/p_0}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}.
\end{aligned}
$$

Since the operator $[\Pi\vec{b},I_\alpha]$ can be devided into the following parts:

$$[\Pi\vec{b},I_\alpha](f_1^0,f_2^\infty)(x)=\mathrm{J}_{21}+\mathrm{J}_{22}+\mathrm{J}_{23}+\mathrm{J}_{24},$$

where

$$
\begin{aligned}
\mathrm{J}_{21}&:=(b_1(x)-b_{1,2Q})(b_2-b_{2,2Q})I_\alpha(f_1^0,f_2^\infty)(x);\\
\mathrm{J}_{22}&:=(b_1(x)-b_{1,2Q})I_\alpha(f_1^0,(b_2-b_{2,2Q})f_2^\infty)(x);\\
\mathrm{J}_{23}&:=(b_2(x)-b_{2,2Q})I_\alpha((b_1-b_{1,2Q})f_1^0,f_2^\infty)(x);\\
\mathrm{J}_{24}&:=I_\alpha((b_1-b_{1,2Q})f_1^0,(b_2-b_{2,2Q})f_2^\infty)(x).
\end{aligned}
$$

This yields

$$\begin{aligned} \mathrm{J}_2 &\lesssim |Q|^{1/q_0-1/q}\bigg(\int_Q |\mathrm{J}_{21}+\mathrm{J}_{22}+\mathrm{J}_{23}+\mathrm{J}_{24}|^q dx\bigg)^{1/q} \\ &\lesssim \|b_1\|_{\mathrm{BMO}}\|b_2\|_{\mathrm{BMO}}\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}. \end{aligned}$$

Combining all the estimates for terms $\mathrm{J}_1, \mathrm{J}_2, \mathrm{J}_3, \mathrm{J}_4$, we get

$$\|[\Pi\vec{b}, I_\alpha](f_1,f_2)\|_{M^{q_0}_q} \lesssim \|b_1\|_{\mathrm{BMO}}\|b_2\|_{\mathrm{BMO}}\|f_1,f_2\|_{\mathcal{M}^{p_0}_{\vec{P}}}.$$

Proof of (2). Let $z_0 \in \mathbb{R}^n$ such that $|(z_0,z_0)| > 2\sqrt{n}$. Take $\mathbb{B} = B\big((z_0,z_0),\sqrt{2n}\big) \subset \mathbb{R}^{2n}$. Since $O \notin \mathbb{B}$, then we can express $(|y_1|^2+|y_2|^2)^{n-\alpha/2}$ as an absolutely convergent Fourier series of the form

$$(|y_1|^2+|y_2|^2)^{n-\alpha/2} = \sum_j a_j e^{iv_j\cdot(y_1,y_2)}, \quad (y_1,y_2)\in B,$$

where $\sum_j |a_j| < \infty$ and we do not care about the vectors $v_j \in \mathbb{R}^{2n}$, but we will at times express them as $v_j = (v_j^1, v_j^2) \in \mathbb{R}^n \times \mathbb{R}^n$.

Let $Q = Q(x_0,r)$ be any arbitrary cube in $\mathbb{R}^n$. Set $\tilde{z} = x_0 + rz_0$ and take $Q' = Q(\tilde{z},r) \subset \mathbb{R}^n$. So for any $x \in Q$ and $y_1, y_2 \in Q'$, we have

$$\Big|\frac{x-y_1}{r} - z_0\Big| \le \Big|\frac{x-x_0}{r}\Big| + \Big|\frac{y_1-\tilde{z}}{r}\Big| \le \sqrt{n}, \quad \Big|\frac{x-y_2}{r} - z_0\Big| \le \Big|\frac{x-x_0}{r}\Big| + \Big|\frac{y_2-\tilde{z}}{r}\Big| \le \sqrt{n},$$

which implies that

$$\bigg(\Big|\frac{x-y_1}{r} - z_0\Big|^2 + \Big|\frac{x-y_2}{r} - z_0\Big|^2\bigg)^{1/2} \le \sqrt{2n};$$

that is, $(\frac{x-y_1}{r}, \frac{x-y_2}{r}) \in \mathbb{B}$.

Let $s(x) = \overline{\mathrm{sgn}(\int_{Q'}(b(x)-b(y))dy)}$. We have the following estimate,

$$\begin{aligned} &\frac{1}{|Q|}\int_Q |b(x)-b_{Q'}|^2 dx \\ &\lesssim \frac{1}{|Q|}\int_Q s^2(x)(b(x)-b_{Q'})^2 dx \\ &\lesssim \frac{1}{|Q||Q'|^2}\int_Q\int_{Q'}\int_{Q'} s^2(x)\big(b(x)-b(y_1)\big)\big(b(x)-b(y_2)\big)dy_1dy_2dx \\ &\lesssim \frac{r^{2n-\alpha}}{|Q|^3}\int_Q\int_{Q'}\int_{Q'} \frac{s^2(x)\big(b(x)-b(y_1)\big(b(x)-b(y_2)\big)\big)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} \sum_j a_j e^{iv_j\cdot(\frac{x-y_1}{r},\frac{x-y_2}{r})}dy_1dy_2dx. \end{aligned}$$

Setting

$$g_j(y_1) = e^{-\frac{i}{r}v_j^1\cdot y_1}\chi_{Q'}(y_1),$$

$$h_j(y_2) = e^{-\frac{i}{r}v_j^2\cdot y_2}\chi_{Q'}(y_2),$$
$$m_j(x) = e^{\frac{i}{r}v_j\cdot(x,x)}\chi_Q(x)s^2(x).$$

We have

$$\begin{aligned}
&\frac{1}{|Q|}\int_Q |b(x)-b_Q|^2dx \lesssim \frac{1}{|Q|}\int_Q |b(x)-b_{Q'}|^2dx\\
&\lesssim \frac{r^{2n-\alpha}\delta^{-2n+\alpha}}{|Q|^3}\sum_j |a_j|\int_Q \big|[\Pi\vec{b},I_\alpha](g_j,h_j)(x)m_j(x)\big|dx\\
&\lesssim \frac{r^{2n-\alpha}}{|Q|^{2+1/q}}\sum_j |a_j|\bigg(\int_Q \big|[\Pi\vec{b},I_\alpha](g_j,h_j)(x)\big|^q dx\bigg)^{1/q}\\
&\lesssim r^{-\alpha-n/q}\sum_j |a_j|\|[\Pi\vec{b},I_\alpha](g_j,h_j)\|_{M_q^{q_0}}\\
&\lesssim \|[\Pi\vec{b},I_\alpha]\|_{\mathcal{M}_{\vec{P}}^{p_0}\to M_q^{q_0}}\sum_j |a_j|.
\end{aligned}$$

The desired result follows from here. □

As mentioned in the introduction, CMO is the closure in BMO of the space of C^∞ functions with compact support. In [37], it was shown that CMO can be characterized in the following way.

Lemma 3.4. *([37]) Let $b\in$ BMO. Then b is in* CMO *if and only if*

$$\lim_{a\to 0}\sup_{|Q|=a}\frac{1}{|Q|}\int_Q |b(x)-b_Q|dx = 0; \tag{3.17}$$

$$\lim_{a\to \infty}\sup_{|Q|=a}\frac{1}{|Q|}\int_Q |b(x)-b_Q|dx = 0; \tag{3.18}$$

$$\lim_{|y|\to 0}\frac{1}{|Q|}\int_Q |b(x+y)-b_Q|dx = 0,\ \ for\ each\ Q. \tag{3.19}$$

Lemma 3.5. *Support that $b\in$ BMO with $\|b\|_*=1$. If for some $0<\epsilon<1$ and a cube Q with its center at x_Q and r_Q, b is not a constant on cube Q and satisfies*

$$\frac{1}{|Q|}\int_Q |b(y)-b_Q|dy > \epsilon^{1/2},$$

then for the function $f_i(i=1,2)$ defined by

$$f_i(y_i) = |Q|^{(\lambda_i-1)/(p_i)}\big(sgn(b(y_i)-b_Q)-c_0\big)\chi_Q(y_i), \tag{3.20}$$

where $c_0 = |Q|^{-1}\int_Q sgn\big(b(y)-b_Q\big)dy_i$ *and* $\frac{\lambda_1}{p_1}+\frac{\lambda_2}{p_2}=\frac{1}{p}-\frac{1}{p_0}$ *with* $0<\lambda_i<1$ *for* $i=1,2$. *There exists constants* $\gamma_1,\gamma_2,\gamma_3$ *satisfying* $\gamma_2>\gamma_1>2$ *and* $\gamma_3>0$, *such that*

$$|Q|^{\frac{1}{q_0}-\frac{1}{q}}\bigg(\int_{\gamma_1 r_Q<|x-x_Q|<\gamma_2 r_Q}\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\bigg)^{1/q}\geq\gamma_3, \tag{3.21}$$

$$|Q|^{\frac{1}{q_0}-\frac{1}{q}}\bigg(\int_{|x-x_Q|>\gamma_2 r_Q}\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\bigg)^{1/q}\leq\frac{\gamma_3}{4}. \tag{3.22}$$

Moreover, there exists a constant $0<\beta<<\gamma_2$ *depending only on* p_1,p_2,n *such that for all measurable subsets* $E\subset\big\{x:\gamma_1 r_Q<|x-x_Q|<\gamma_2 r_Q\big\}$ *satisfying* $\frac{|E|}{|Q|}<\beta^n$, *we have*

$$|Q|^{\frac{1}{q_0}-\frac{1}{q}}\bigg(\int_{E}\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\bigg)^{1/q}\leq\frac{\gamma_3}{4}. \tag{3.23}$$

Proof. Since $\int_Q\big(b(y)-b_Q\big)dy=0$, it is easy to check that f_i satisfies

$$\begin{gathered}\mathrm{supp} f_i\subset Q,\\ f_i(y_i)(b(y)-b_Q)\geq 0,\\ \int f_i(y_i)dy_i=0,\\ |f_i(y_i)|\leq 2|Q|^{(\lambda_i-1)/p_i},\\ \int\big(b(y)-b_Q\big)f_i(y)dy=|Q|^{(\lambda_i-1)/p_i}\int_Q|b(y_i)-b_Q|dy.\end{gathered}$$

Moreover, it is easy to see that $\|(f_1,f_2)\|_{\mathcal{M}^{p_0}_p}\leq C$. For a cube Q with center x_Q and $x\in(2\sqrt{n}Q)^c$, the following point-wise estimates hold:

$$|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)|\lesssim|Q|^{2-\frac{1}{p_0}}|x-x_Q|^{-2n+\alpha}, \tag{3.24}$$

$$|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)|\gtrsim\epsilon|Q|^{2-\frac{1}{p_0}}|x-x_Q|^{-2n+\alpha}, \tag{3.25}$$

$$|I_\alpha\big((b-b_Q)f_1,f_2\big)(x)|\lesssim|Q|^{2-\frac{1}{p_0}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}, \tag{3.26}$$

$$|I_\alpha\big(f_1,(b-b_Q)f_2\big)(x)|\lesssim|Q|^{2-\frac{1}{p_0}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}, \tag{3.27}$$

$$|I_\alpha\big(f_1,f_2\big)(x)|\lesssim|Q|^{2-\frac{1}{p_0}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}, \tag{3.28}$$

where f_i as above and the constants involved are independent of b, f_i and ϵ.

To prove (3.24), from the fact that $\|b\|_*=1$ and $x\in(2\sqrt{n}Q)^c$, we have

$$|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)|$$

$$
\begin{aligned}
&= \Big| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)(b(y_2)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \Big| \\
&\lesssim |x-x_Q|^{-2n+\alpha} \prod_{i=1}^2 \int_Q (b(y_i)-b_Q)f_i(y_i)dy_i \\
&\lesssim |Q|^{\frac{\lambda_1-1}{p_1}+\frac{\lambda_2-1}{p_2}} |x-x_Q|^{-2n+\alpha} \prod_{i=1}^2 \int_Q |b(y_i)-b_Q|dy_i \\
&\lesssim |Q|^{2+\frac{\lambda_1}{p_1}+\frac{\lambda_2}{p_2}-\frac{1}{p_1}-\frac{1}{p_2}} |x-x_Q|^{-2n+\alpha} \\
&\lesssim |Q|^{2-\frac{1}{p_0}} |x-x_Q|^{-2n+\alpha}.
\end{aligned}
$$

For (3.25), using that $\big(b(y_i)-b_Q\big)f_i(y_i)\geq 0$, we can compute

$$
\begin{aligned}
&|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)| \\
&= \Big| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)(b(y_2)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \Big| \\
&\gtrsim |x-x_Q|^{-2n+\alpha} \prod_{i=1}^2 \Big| \int_Q \big(b(y_i)-b_Q\big)f_i(y_i)dy_i \Big| \\
&= |x-x_Q|^{-2n+\alpha} \prod_{i=1}^2 |Q|^{\frac{\lambda_i-1}{p_i}} \int_Q \big|b(y_i)-b_Q\big|dy_i \\
&\gtrsim \epsilon |Q|^{2-\frac{1}{p_0}} |x-x_Q|^{-2n+\alpha}.
\end{aligned}
$$

For (3.26), applying the fact $|f_2(y_2)|\leq 2|Q|^{(\lambda_2-1)/p_2}$ and $\int_Q f_2(y_2)dy_2=0$, we can also estimate for any $y'\in Q$,

$$
\begin{aligned}
&|I_\alpha\big((b-b_Q)f_1,f_2\big)(x)| \\
&= \Big| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \Big| \\
&= \Big| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \\
&\qquad - \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2'|^2\big)^{n-\alpha/2}} dy_1dy_2 \Big| \\
&\lesssim |Q|^{\frac{\lambda_2-1}{p_2}} \int_Q\int_Q \frac{|y_2-y_2'|(b(y_1)-b_Q)f_1(y_1)}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}} dy_1dy_2
\end{aligned}
$$

$$\lesssim |Q|^{\frac{\lambda_1-1}{p_1}+\frac{\lambda_2}{p_2}+\frac{1}{p_2'}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}\int_Q |b(y_1)-b_Q|dy_1$$
$$\lesssim |Q|^{2-\frac{1}{p_0}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}.$$

It is easy to see that $|I_\alpha\big((b-b_Q)f_1,f_2\big)(x)|=|I_\alpha\big(f_1,(b-b_Q)f_2\big)(x)|$, then (3.27) holds. Finally, using that f_1 has mean zero we obtain (3.28) as follows.

$$\begin{aligned}
&|I_\alpha\big(f_1,f_2\big)(x)|\\
&=\Big|\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}}-\frac{f_1(y_1)f_2(y_2)}{\big(|x-y_1'|^2+|x-y_2|^2\big)^{n-\alpha/2}}dy_1dy_2\Big|\\
&\lesssim\int_Q\int_Q\frac{|y_1-y_1'||f_1(y_1)||f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}}dy_1dy_2\\
&\lesssim |Q|^{2-\frac{1}{p_0}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}.
\end{aligned}$$

Now, we give the proofs of (3.21)-(3.23). Taking $\nu>16$, by (3.26) we obtain

$$\begin{aligned}
&\Big(\int_{|x-x_Q|>\nu r_Q}\big|(b(x)-b_Q)I_\alpha((b-b_Q)f_1,f_2)(x)\big|^q dx\Big)^{\frac{1}{q}}\\
&\lesssim |Q|^{2-\frac{1}{p_0}+\frac{1}{n}}\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}\Big(\int_{2^sr_Q<|x-x_Q|<2^{s+1}r_Q}\frac{|b(x)-b_Q|^q}{|x-x_Q|^{q(2n-\alpha+1)}}dx\Big)^{\frac{1}{q}}\\
&\lesssim |Q|^{1/q_0-1/q}\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}2^{-s(2n-\alpha+1)}\Big(\frac{1}{|2^{s+1}Q|}\int_{2^sr_Q<|x-x_Q|<2^{s+1}r_Q}|b(x)-b_Q|^q dx\Big)^{\frac{1}{q}}\\
&\lesssim |Q|^{1/q-1/q_0}\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}s2^{-s(2n-\alpha+1-\frac{n}{q})}\\
&\lesssim |Q|^{1/q-1/q_0}\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}2^{-s(2n-\alpha-\frac{n}{q}+\frac{1}{2})}\\
&\lesssim |Q|^{1/q-1/q_0}\nu^{-(2n-\alpha-\frac{n}{q}+\frac{1}{2})},
\end{aligned}$$

where we have used that $s\le 2^{s/2}$ for $4\le\lfloor\log_2\nu\rfloor\le s$.

Similarly, we also have

$$\Big(\int_{|x-x_Q|>\nu r_Q}\big|(b(x)-b_Q)I_\alpha(f_1,(b-b_Q)f_2)(x)\big|^q dx\Big)^{\frac{1}{q}}\lesssim |Q|^{1/q_0-1/q}\nu^{-2n-\alpha+\frac{1}{2}-\frac{n}{q}},$$

$$\Big(\int_{|x-x_Q|>\nu r_Q}\big|(b(x)-b_Q)^2I_\alpha(f_1,f_2)(x)\big|^q dx\Big)^{\frac{1}{q}}\lesssim |Q|^{1/q_0-1/q}\nu^{-2n-\alpha+\frac{1}{2}-\frac{n}{q}}.$$

Then for $\mu > \nu$, using (3.24), (3.25) and the estimates above, we get

$$
\begin{aligned}
&|Q|^{1/q_0-1/q}\int_{\nu r_Q<|x-x_Q|<\mu r_Q}\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\\
&\geq C|Q|^{1/q_0-1/q}\bigg(\int_{\nu r_Q<|x-x_Q|<\mu r_Q}\big|I_\alpha((b-b_Q)f_1,(b-b_Q)f_2)(x)\big|^q dx\bigg)^{1/q}\\
&\quad -C|Q|^{1/q_0-1/q}\bigg(\int_{\nu r_Q<|x-x_Q|}\big|(b(x)-b_Q)I_\alpha((b-b_Q)f_1,f_2)(x)\big|^q dx\bigg)^{1/q}\\
&\quad -C|Q|^{1/q_0-1/q}\bigg(\int_{\nu r_Q<|x-x_Q|}\big|(b(x)-b_Q)I_\alpha(f_1,(b-b_Q)f_2)(x)\big|^q dx\bigg)^{1/q}\\
&\quad -C|Q|^{1/q_0-1/q}\bigg(\int_{\nu r_Q<|x-x_Q|}\big|(b(x)-b_Q)^2I_\alpha(f_1,f_2)(x)\big|^q dx\bigg)^{1/q}\\
&\geq C\epsilon|Q|^{2-\frac{\alpha}{n}-\frac{1}{q}}\bigg(\int_{\nu r_Q<|x-x_Q|<\mu r_Q}|x-x_Q|^{q(-2n+\alpha)}dx\bigg)^{1/q}-C\nu^{-2n+\alpha+\frac{n}{q}-\frac{1}{2}}\\
&\geq C\epsilon\Big(\nu^{-2nq+n+\alpha q}-\mu^{-2nq+n+\alpha q}\Big)^{\frac{1}{q}}-C\nu^{-2n+\alpha+\frac{n}{q}-\frac{1}{2}}.
\end{aligned}
$$

We can select γ_1,γ_2 in place of ν,μ with $\gamma_2 >> \gamma_1$, then (3.21) and (3.22) are verified for some $\gamma_3 > 0$.

We now verified (3.23). Let $E \subset \big\{\gamma_1 r_Q < |x-x_Q| < \gamma_2 r_Q\big\}$ be an arbitrary measurable set. It follows from Minkowski inequality that

$$
\begin{aligned}
&|Q|^{1/q_0-1/q}\bigg(\int_E\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\bigg)^{\frac{1}{q}}\\
&\lesssim |Q|^{2-\alpha/n-1/q}\bigg(\int_E|x-x_Q|^{-q(2n-\alpha)}dx\bigg)^{\frac{1}{q}}+|Q|^{2-\alpha/n-1/q+\frac{1}{n}}\bigg(\int_E\frac{|b(x)-b_Q|^q}{|x-x_Q|^{q(2n-\alpha+1)}}dx\bigg)^{\frac{1}{q}}\\
&\qquad +|Q|^{2-\alpha/n-1/q+\frac{1}{n}}\bigg(\int_E\frac{|b(x)-b_Q|^{2q}}{|x-x_Q|^{q(2n-\alpha+1)}}dx\bigg)^{\frac{1}{q}}\\
&\lesssim \bigg(\frac{|E|^{1/q}}{|Q|^{1/q}}+\Big(\frac{1}{|Q|}\int_E|b(x)-b_Q|^q dx\Big)^{\frac{1}{q}}+\Big(\frac{1}{|Q|}\int_E|b(x)-b_Q|^{2q}dx\Big)^{\frac{1}{q}}\bigg)\\
&\lesssim \frac{|E|^{\frac{1}{2q}}}{|Q|^{\frac{1}{2q}}}\bigg(1+\log\Big(\frac{\tilde{C}|Q|}{|E|}\Big)\bigg)^{\frac{1+\lfloor 2q\rfloor}{2q}}.
\end{aligned}
$$

The last inequality can be obtained by [12, P.309] taking $0 < \beta < \min\{\tilde{C}^{1/n},\gamma_2\}$ and sufficiently small so that (3.23) holds. □

In order to prove Theorem 2.1 and 2.2, we need the characterization that a subset of $M_p^{p_0}$ is a strong pre-compact set.

Lemma 3.6. *[14] Let $1 < p \le p_0 < \infty$. Suppose that the subset $\mathfrak{F} \subset M_p^{p_0}$ satisfies the following conditions:*

(i) norm boundedness uniformly

$$\sup_{f\in\mathfrak{F}} \|f\|_{M_p^{p_0}} < \infty;$$

(ii) control uniformly away from the origin

$$\lim_{A\to\infty} \|f\chi_{E_A}\|_{M_p^{p_0}} = 0 \ uniformly \ in \ f \in \mathfrak{F}, where \ E_A = \{x \in \mathbb{R}^n : |x| > A\};$$

(iii) translation continuity uniformly

$$\lim_{y\to 0} \|f(\cdot + y) - f(\cdot)\|_{M_p^{p_0}} = 0 \ uniformly \ in \ f \in \mathfrak{F};$$

then $\mathfrak{F}$ is pre-compact in $M_p^{p_0}$.

4. Proof of Theorem 2.1 and Theorem 2.2

Proof of Theorem 2.1. We need only to show the set $\mathfrak{F} := \{[\Pi\vec{b}, T](f_1, f_2) : \|(f_1, f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}} \le 1\}$ is strong pre-compact in $M_p^{p_0}$ when $b_1, b_2 \in C_c^\infty$. By Lemma 3.6, we need to verify the conditions $(i), (ii)$ and (iii) hold uniformly in $\mathfrak{F}$ for $b_1, b_2 \in C_c^\infty$.

It is easy to verify that $\mathfrak{F}$ satisfies the condition (i) by Lemma 3.1.

As for condition (ii), suppose that supp $b_1, b_2 \subset \{x \in \mathbb{R}^n : |x| \le \beta\}$ with $\beta > 1$. For any $A > 2\beta$,

$$|Q|^{1/p_0-1/p}\bigg(\int_Q |[\Pi\vec{b}, T](f_1, f_2)(x)|^p \chi_{E_A}(x)dx\bigg)^{1/p} \to 0, (A \to +\infty).$$

In fact, for any $x \in E_A$, $|x| \le |x - y_1| + |y_1| \le |x - y_1| + |x - y_2|$, then

$$\begin{aligned} |[\Pi\vec{b}, T](f_1, f_2)(x)| &\lesssim \|b_1\|_\infty \|b_2\|_\infty \int_{|y_1|\le\beta}\int_{|y_2|\le\beta} \frac{|f_1(y_1)f_2(y_2)|}{(|x - y_1| + |x - y_2|)^{2n}} dy_1 dy_2 \\ &\lesssim \frac{1}{|x|^{2n}} \int_{|y_1|\le\beta}\int_{|y_2|\le\beta} |f_1(y_1)f_2(y_2)|dy_1 dy_2 \\ &\lesssim \beta^{2n-n/p_0}|x|^{-2n}\|(f_1, f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}}. \end{aligned}$$

From $L^{p_0} \subset M_q^{p_0}$, it follows that for any cube Q

$$|Q|^{1/p_0-1/p}\bigg(\int_Q |[\Pi\vec{b}, T](f_1, f_2)(x)|^p \chi_{E_A}(x)dx\bigg)^{1/p}$$

$$\lesssim \Big(\int_{|x|>A} \beta^{2n-n/p_0}|x|^{-2n}dx\Big)^{1/p_0}$$
$$\lesssim \Big|\frac{\beta}{A}\Big|^{2n-n/p_0}.$$

Thus, the inequality above tends to zero as $A \to \infty$.

Finally, it remains to prove condition (iii). We need to show that for any $0 < \epsilon < 1$, if $|t|$ is sufficiently small depending ϵ, then

$$\|[\Pi\vec{b}, T](f_1, f_2)(\cdot + t) - [\Pi\vec{b}, T](f_1, f_2)(\cdot)\|_{M^{p_0}_p} \lesssim \epsilon.$$

To do this, we break

$$[\Pi\vec{b}, T](f_1, f_2)(x) - [\Pi\vec{b}, T](f_1, f2)(x + t)$$

into a sum of four terms

$$\mathrm{II}_1 + \mathrm{II}_2 + \mathrm{II}_3 + \mathrm{II}_4$$

with

$$\mathrm{II}_1 := \iint_\Omega K(x, y_1, y_2)(b_1(x + t) - b_1(x))(b_2(y_2) - b_2(x))f_1(y_1)f_2(y_2)dy_1dy_2;$$
$$\mathrm{II}_2 := \iint_\Omega \Big(K(x, y_1, y_2)(b_2(y_2) - b_2(x)) - K(x + t, y_1, y_2)(b_2(y_2) - b_2(x + t))\Big)$$
$$\times(b_1(y_1) - b_1(x + t))f_1(y_1)f_2(y_2)dy_1dy_2;$$
$$\mathrm{II}_3 := \iint_{\Omega^c} K(x, y_1, y_2)(b_1(y_1) - b_1(x))(b_2(y_2) - b_2(x))f_1(y_1)f_2(y_2)dy_1dy_2;$$
$$\mathrm{II}_4 := \iint_{\Omega^c} K(x + t, y_1, y_2)(b_1(y_1) - b_1(x + t))(b_2(x + t) - b_2(y_2))f_1(y_1)f_2(y_2)dy_1dy_2,$$

where $\Omega = \{(y_1, y_2) : |x - y_1| + |x - y_2| > \delta := \epsilon^{-1}|t|\}$ and $\Omega^c = \mathbb{R}^{2n}\backslash\Omega$.

For II_1, we can compute

$$\mathrm{II}_1(\mathrm{x}, \mathrm{t}) \lesssim |t|\|\nabla b_1\|_\infty\big(T_*(f_1, b_2f_2)(x) + |b_2(x)|T_*(f_1, f_2)(x)\big).$$

By Corollary 3.1 and $b_2 \in L^\infty$, we obtain

$$\|\mathrm{II}_1\|_{M^{p_0}_p} \lesssim |t|\bigg(\|(f_1, b_2f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}} + \|b_2\|_\infty\|(f_1, f_2)\|_{\mathcal{M}^{p_0}_{\vec{P}}}\bigg) \lesssim |t|.$$

To deal with the II_2 term, we write II_2 as a sum of three terms, $\mathrm{II}_{21} + \mathrm{II}_{22} + \mathrm{II}_{23}$, where

$$\mathrm{II}_{21} = \iint_\Omega \Big(K(x, y_1, y_2) - K(x + t, y_1, y_2)\Big)(b_1(y_1) - b_1(x + t))f_1(y_1)b_2(y_2)f_2(y_2)dy_1dy_2;$$
$$\mathrm{II}_{22} = b_2(x + t)\iint_\Omega \Big(K(x, y_1, y_2) - K(x + t, y_1, y_2)\Big)(b_1(y_1) - b_1(x))f_1(y_1)f_2(y_2)dy_1dy_2.$$

$$\mathrm{II}_{23}=\iint_{\Omega}K(x,y_1,y_2)(b_1(y_1)-b_1(x+t))(b_2(x+t)-b_2(x))f_1(y_1)f_2(y_2)dy_1dy_2.$$

By the regularity condition of function K, we have

$$\begin{aligned}\mathrm{II}_{21} &\lesssim \|b_1\|_\infty\|b_2\|_\infty\iint_{\Omega}\frac{|t|^\gamma|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n+\gamma}}dy_1dy_2\\ &\lesssim |t|^\gamma\sum_{k=1}^{\infty}\iint_{2^{k-1}\delta\le|x-y_1|+|x-y_2|<2^k\delta}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n+\gamma}}dy_1dy_2\\ &\lesssim |t|^\gamma\delta^{-\gamma}\mathcal{M}(f_1,f_2)(x),\end{aligned}$$

where the bilinear maximal operator $\mathcal{M}$ is defined by Lerner et al.[30], which is used to obtain a precise control on multinear singular integral operators. The bilinear maximal operator $\mathcal{M}$ is defined by

$$\mathcal{M}(f_1,f_2)(x)=\sup_{Q\ni x}\prod_{i=1}^{2}\frac{1}{|Q|}\int_Q|f_i(y_i)|dy_i.$$

By the boundedness of $\mathcal{M}$ from $\mathcal{M}^{p_0}_{\vec{P}}$ to $M^{p_0}_p$ (see [25]), we obtain

$$\|\mathrm{II}_{21}\|_{M^{p_0}_p}\lesssim|t|^\gamma\delta^{-\gamma}\|\mathcal{M}(f_1,f_2)\|_{M^{p_0}_p}\lesssim|t|^\gamma\delta^{-\gamma}.$$

The term II_{22} is estimated using the same methods as in the proof for II_{21}, and the term II_{23} is same as II_1. Then

$$\|\mathrm{II}_2\|_{M^{p_0}_p}\lesssim|t|+|t|^\gamma\delta^{-\gamma}.$$

Note that

$$\begin{aligned}\mathrm{II}_3 &\lesssim \|\nabla b_1\|_{L^\infty}\|\nabla b_2\|_{L^\infty}\iint_{\Omega^c}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n-2}}dy_1dy_2\\ &\lesssim \sum_{k=1}^{\infty}\iint_{2^{-k}\delta<|x-y_1|+|x-y_2|\le2^{-k+1}\delta}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|+|x-y_2|)^{2n-2}}dy_1dy_2\\ &\lesssim \delta^2\mathcal{M}(f_1,f_2)(x),\end{aligned}$$

which gives that

$$\|\mathrm{II}_3\|_{M^{p_0}_p}\lesssim\delta^2.$$

Finally, for the last term we proceed in an analogous manner, by replacing x with $x+t$ and the region of integration $\{(y_1,y_2):|x-y_1|+|x-y_2|<\delta\}$ with the larger one $\{(y_1,y_2):|x+t-y_1|+|x+t-y_2|<\delta+2|t|\}$. Thus,

$$\|\mathrm{II}_4\|_{M^{p_0}_p}\lesssim(\delta+2|t|)^2.$$

For any $0 < \epsilon < 1$, there exists a constant $t_0 = \epsilon^2$, for any $|t| < t_0$,

$$\|[\Pi\vec{b}, T](f_1, f_2)(\cdot + t) - [\Pi\vec{b}, T](f_1, f_2)(\cdot)\|_{M_p^{p_0}} \lesssim |t| + \epsilon^\gamma + \frac{|t|}{\epsilon} + (\frac{1}{\epsilon} + 2)^2|t|^2 \lesssim \epsilon^\gamma.$$

We prove that condition (iii) holds for $[\Pi\vec{b}, T](f_1, f_2)$ uniformly in $\mathfrak{F}$ and Theorem 2.1 follows.

Proof of Theorem 2.2. We need only to verify the conditions $(i), (ii)$ and (iii) hold uniformly in $\mathcal{G}$ for $b_1, b_2 \in C_c^\infty$, where

$$\mathcal{G} = \big\{[\Pi\vec{b}, I_\alpha](f_1, f_2) : \|(f_1, f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}} \leq 1\big\}.$$

By Lemma 3.3, we have $\mathcal{G}$ is uniformly bounded.

For the condition (ii), suppose that supp $b_1, b_2 \subset \{x \in \mathbb{R}^n : |x| \leq \beta\}$ with $\beta > 1$ and let $A \geq 2\beta$. Then for any $|x| > A$ and $|y_1|, |y_2| \leq \beta$, we have $|x - y_1|^2 + |x - y_2|^2 \gtrsim |x|^2$. Thus, for any cube Q,

$$\begin{aligned}
&|Q|^{1/q_0 - 1/q}\bigg(\int_Q |[\Pi\vec{b}, I_\alpha](f_1, f_2)(x)|^q \chi_{E_A}(x)dx\bigg)^{1/q} \\
&\lesssim |Q|^{1/q_0 - 1/q}\bigg(\int_Q \bigg(\int_{|y_1|\lesssim\beta}\int_{|y_2|\lesssim\beta} \frac{|f_1(y_1)f_2(y_2)|dy_1dy_2}{(|x - y_1|^2 + |x - y_2|^2)^{n-\alpha/2}}\bigg)^q \chi_{E_A}(x)dx\bigg)^{1/q} \\
&\lesssim |Q|^{1/q_0 - 1/q}\bigg(\int_Q |x|^{\alpha q - 2nq}\beta^{2nq - nq/q_0}\chi_{E_A}(x)dx\bigg)^{1/q} \|(f_1, f_2)\|_{\mathcal{M}_{\vec{P}}^{p_0}} \\
&\lesssim \beta^{2n - n/q_0}\bigg(\int_{|x|>A} |x|^{\alpha q_0 - 2nq_0}dx\bigg)^{1/q_0} \\
&\lesssim (\frac{\beta}{A})^{2n - n/q_0}.
\end{aligned}$$

Thus, (b) holds by letting $A \to \infty$.

To prove the uniform continuity of $\mathcal{G}$, we must see that

$$\lim_{t\to 0} \|[\Pi\vec{b}, I_\alpha](f_1, f_2)(\cdot + t) - [\Pi\vec{b}, I_\alpha](f_1, f_2)(\cdot)\|_{M_q^{q_0}} = 0.$$

To deal with compactness of fractional integral operators, we find it convenient to use smooth truncations of I_α. The operator I_α^δ is defined by a smooth kernel $K^\delta(x, y_1, y_2)$ such that

$$K^\delta(x, y_1, y_2) = \frac{1}{(|x - y_1|^2 + |x - y_2|^2)^{n-\alpha/2}}$$

for $|x - y_1| + |x - y_2| > 2\delta$;

$$K^\delta(x, y_1, y_2) = 0$$

for $|x-y_1|+|x-y_2|\le\delta$; and

$$|\partial^\gamma K^\delta(x,y_1,y_2)|\lesssim\frac{1}{(|x-y_1|+|x-y_2|)^{2n-\alpha+\gamma}}$$

for all (x,y_1,y_2) and all-multi-indexes with $|\gamma|\le 1$.

Then, we need only to show that

$$\lim_{t\to 0}\|[\Pi\vec{b},I_\alpha^\delta](f_1,f_2)(\cdot+t)-[\Pi\vec{b},I_\alpha^\delta](f_1,f_2)(\cdot)\|_{M_q^{q_0}}=0, \tag{4.1}$$

where $\delta=|t|^{1/2}$. In fact, for any $x\in\mathbb{R}^n$

$$\begin{aligned}
&\Big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)-[\Pi\vec{b},I_\alpha^\delta](f_1,f_2)(x)\Big|\\
&\le\iint_{|x-y_1|+|x-y_2|\le 2\delta}\frac{|b_1(x)-b_1(y_1)||b_2(x)-b_2(y_2)||f_1(y_1)||f_2(y_2)|}{(|x-y_1|^2+|x-y_2|^2)^{n-\alpha/2}}dy_1dy_2\\
&\lesssim\|\nabla b_1\|_\infty\|\nabla b_2\|_\infty\iint_{|x-y_1|+|x-y_2|\le 2\delta}\frac{|f_1(y_1)||f_2(y_2)|}{(|x-y_1|^2+|x-y_2|^2)^{n-\alpha/2-1}}dy_1dy_2\\
&\lesssim\sum_{k=0}^{\infty}(2^{-k}\delta)^{\alpha+2-2n}\iint_{2^{-k+1}\delta<|x-y_1|+|x-y_2|\le 2^k\delta}|f_1(y_1)||f_2(y_2)|dy_1dy_2\\
&\lesssim\sum_{k=0}^{\infty}(2^{-k}\delta)^2\mathcal{M}_\alpha(f_1,f_2)(x)\\
&\lesssim\delta^2 I_\alpha(|f_1|,|f_2|)(x).
\end{aligned}$$

Set $\delta^2=|t|$, which gives that

$$\|[\Pi\vec{b},I_\alpha](f_1,f_2)-[\Pi\vec{b},I_\alpha^\delta](f_1,f_2)\|_{M_q^{q_0}}\lesssim|t|.$$

To prove (4.1), we write

$$\begin{aligned}
&[\Pi\vec{b},I_\alpha^\delta](f_1,f_2)(x)-[\Pi\vec{b},I_\alpha^\delta](f_1,f_2)(x+t)\\
&=\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\big(b_1(x)-b_1(y_1)\big)\big(b_2(x)-b(y_2)\big)K^\delta(x,y_1,y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&\quad-\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\big(b_1(x+t)-b(y_1)\big)\big(b_2(x+t)-b(y_2)\big)K^\delta(x+t,y_1,y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&=\big(b_1(x)-b_1(x+t)\big)\big(b_2(x)-b_2(x+t)\big)\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}K^\delta(x,y_1,y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&\quad+\big(b_1(x)-b_1(x+t)\big)\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\big(b_2(x+t)-b(y_2)\big)K^\delta(x,y_1,y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&\quad+\big(b_2(x)-b_2(x+t)\big)\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\big(b_1(x+t)-b(y_1)\big)K^\delta(x,y_1,y_2)f_1(y_1)f_2(y_2)dy_1dy_2
\end{aligned}$$

$$+\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\big(b_1(x+t)-b(y_1)\big)\big(b_2(x+t)-b(y_2)\big)$$
$$\times\big(K^\delta(x,y_1,y_2)-K^\delta(x+t,y_1,y_2)\big)f_1(y_1)f_2(y_2)dy_1dy_2$$
$$=\mathrm{JJ}_1+\mathrm{JJ}_2+\mathrm{JJ}_3+\mathrm{JJ}_4.$$

For JJ_1, we simply have

$$|\mathrm{JJ}_1|\lesssim|t|^2\|\nabla b_1\|_\infty\|\nabla b_2\|_\infty I_\alpha(|f_1|,|f_2|)(x),$$

which implies that

$$\|\mathrm{JJ}_1\|_{M^{q_0}_q}\lesssim|t|^2.$$

Similarly, we also have that for $j=2,3$

$$|\mathrm{JJ_j}|\lesssim|t|\|\nabla b_1\|_\infty\|b_2\|_\infty I_\alpha(|f_1|,|f_2|)(x).$$

We now give the estimate for JJ_4. We may assume that t small enough such that $|t|\in(0,\frac{\delta}{4})$. If $|x-y_1|+|x-y_2|\le\delta/2$ we have $|x+t-y_1|+|x+t-y_2|\le\delta/2+2|t|\le\delta$, thus

$$K^\delta(x+t,y_1,y_2)-K^\delta(x,y_1,y_2)=0.$$

This gives us that

$$\begin{aligned}
\big|\mathrm{JJ}_4\big| &\lesssim |t|\|b_1\|_\infty\|b_2\|_\infty\iint_{|x-y_1|+|x-y_2|>\frac{\delta}{2}}\frac{|f_1(y_1)f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}}dy_1dy_2\\
&\lesssim |t|\sum_{j=0}^{\infty}\iint_{2^{j-1}\delta<|x-y_1|+|x-y_2|<2^j\delta}\frac{|f_1(y_1)f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}}dy_1dy_2\\
&\lesssim |t|\sum_{j=0}^{\infty}(2^j\delta)^{-1}\iint_{2^{j-1}\delta<|x-y_1|+|x-y_2|<2^j\delta}\frac{|f_1(y_1)f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha}}dy_1dy_2\\
&\lesssim \frac{|t|}{\delta}I_\alpha(|f_1|,|f_2|)(x).
\end{aligned}$$

Combining the estimates above and t small enough such that $0<4t<\delta:=|t|^{1/2}$, we have

$$\|[\Pi\vec{b},I^\delta_\alpha](f_1,f_2)(\cdot+t)-[\Pi\vec{b},I^\delta_\alpha](f_1,f_2)(\cdot)\|_{M^{q_0}_q}\lesssim(|t|^2+|t|+\frac{|t|}{\delta})\|I_\alpha(|f_1|,|f_2|)\|_{M^{q_0}_q}\lesssim|t|^{1/2}.$$

Thus, the compactness of $[\Pi\vec{b},I_\alpha]$ on Morrey space is completed.

So it remains to show that if $b_1=b_2$ and $[\Pi\vec{b},I_\alpha]$ is a compact operator from $\mathcal{M}^{p_0}_{\vec{P}}$ to $M^{q_0}_q$, then $b_1,b_2\in\mathrm{CMO}$.

First, Lemma 3.3 implies that $b \in \mathrm{BMO}$. To prove b be an element of CMO, we will adapt some arguments from [12], see also [7], which in turn are based on the original work in [37]. The approach is the following: if one of the conditions Eqs.(3.17)-(3.19) in Lemma 3.4 is failed, we will show that there exist sequences of functions, $\{f_j\}_j$ and $\{g_j\}_j$ uniformly bounded on $\mathcal{M}^{p_0}_{\vec{P}}$, such that $[[\Pi\vec{b}, I_\alpha](f_j, g_j)$ has no convergent subsequence, which contradicts the assumption that $[\Pi\vec{b}, I_\alpha]$ is compact. It gives us that if $[\Pi\vec{b}, I_\alpha]$ is compact, b must satisfy all three conditions; that is $b \in \mathrm{CMO}$.

By Lemma 3.5, it is sufficient to once again repeat the steps preformed in [7] (or[12],[13]) to obtain the desired result and it is left to the reader. □

Acknowledgments We would like to thank the anonymous referee for his/her comments.

References

[1] Adams, D.: A note on Riesz potentials. Duke Math. J. **42**(4), 765-778 (1975)

[2] Adams, D., Xiao, J.: Nonlinear potential analysis on Morrey spaces and their capacities. Indiana Univ. Math. J. **53**(6),1629-1663 (2004)

[3] Bényi, Á., Damián, W., Moen, K., Torres, R.H.: Compactness properties of commutators of bilinear frctional integrals. Math. Z. **280**, 569-582 (2015)

[4] Bényi, Á., Damián, W., Moen, K., Torres, R.H.: Compact bilinear operators: the weighted case. Mich. Math. J. **4**, 39-51 (2015)

[5] Bényi, Á., Oh, T.: Smoothing of commtators for a Hörmander classes of bilinear pseudodifferential operators. J. Fourier Anal. Appl. **20**(2), 282-300 (2014)

[6] Bényi, Á., Torres, R.H.: Compact bilinear operators and commutators. Proc. Amer. Math. Soc. **141**(10), 3609-3621 (2013)

[7] Chaffee, L., Torres, R.H.: Characterization of Compactness of the Commutators of Bilinear Fractional Integral Operators. Potential Anal. **43**, 481-494 (2015)

[8] Chanillo, S.: *A note on commutators*, Indiana Univ. Math. J., 1982(31): 7-16.

[9] Chen, Y., Ding, Y.: Compactness of commutators of singular integrals with variable kernels, (in Chinese). Chin. Ann. Math. **30**(A), 201-212(2009)

[10] Chen, Y., Ding, Y.: Compactness of commutators for Littlewood-Paley operators. Kodai Math. J. **32**, 256-323 (2009)

[11] Chen, Y., Ding, Y.: Compactness of the commutators for parabolic singular integrals. Sci. China (A) **53**, 2633-2648 (2010)

[12] Chen, Y., Ding, Y., Wang, X.: Compactness of commutators of Riesz potential on Morrey spaces. Potential Anal. **30**, 301-313 (2009)

[13] Chen, Y., Ding, Y., Wang, X.: Compactness for commutators of Marcinkiewicz integral in Morrey spaces. Taiwan. J. Math. **15**, 633-658 (2011)

[14] Chen, Y., Ding, Y., Wang, X.: Compactness of commutators for Singular integrals on Morrey spaces. Canad. J. Math. **64**, 257-281(2012)

[15] Christ, M., Journé, J.L.: Polynomial growth estimates for multilinear singular integral operators, Acta Math. **159**,51-80 (1987)

[16] Clop, A., Cruz, V.: Weighted estimates for Beltrami equations. Ann. Acad. Sci. Fenn. Math. **38**, 91-113 (2013)

[17] Coifman, R., Meyer, Y.: On commutators of singular integrals and bilinear singular integrals, Trans. Amer. Math. **212**, 315-331 (1975)

[18] Coifman, R., Meyer, Y.: Commutateurs d'intégrales singulières et opérateurs multilinéaires, Ann. Inst. Fourier (Grenoble) **28**(3), 177-202 (1978)

[19] Coifman, R., Rochberg, R., Weiss, G.: Factorization theorems for Hardy spaces in several variables. *Ann. of Math.*, **103**, 611-635 (1976)

[20] Ding, Y., Mei, T.: Boundedness and compactness for the commutators of bilinear operators on Morrey spaces. Potential Anal. **42**, 717-748 (2015)

[21] Grafakos, L.: On multilinear fractional integrals. Studia Math. **102**, 49-56 (1992)

[22] Grafakos, L., Kalton, N.: Some remarks on multilinear maps and interpolation, Math. Ann. **319**, 151-180 (2001)

[23] Grafakos, L., Torres, R.H.: Multilinear Calderon-Zygmund theory, Adv. Math. **165**(1), 124-164 (2002)

[24] Hanche-Olsen, H., Holden, H.: The Kolmogorov-Riesz compactness theorem. Expo. Math. **28**, 385-394 (2010)

[25] Iida, T., Sato, E., Sawano, Y., Tanaka, H.: Sharp bounds for multilinear fractional integral operators on Morrey type spaces. Positivity **16**, 339-358 (2012)

[26] Kato, T.: Strong solutions of the Navier-Stokes equation in Morrey spaces. Bol. Soc. Brasil. Mat. **22**(2), 127-155 (1992)

[27] Kenig, C., Stein, E.: Multilinear estimates and fractional integration, Math. Res. Lett. **6**, 1-15 (1999)

[28] Krantz, S., Li, S.-Y.: Boundedness and compactness of integral operators on spaces of homogeneous type and applications, I. J. Math. Anal. Appl. **258**, 629-641 (2001)

[29] Krantz, S., Li, S.-Y.: Boundedness and compactness of integral operators on spaces of homogeneous type and applications, II. J. Math. Anal. Appl. **258**, 642-657 (2001)

[30] Lerner, A.K., Ombrosi, S., Pérez, C., Torres, R.H., Trujillo-González, R.: New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory. *Adv. Math.*, **220**, 1222-1264 (2009)

[31] Moen, K.: Weighted inequalities for multilinear fractional integral operators. Collect Math. **60**, 213-238 (2009)

[32] Pérez, C.: Endpoint estimates for commutators of singular integral operators. *J. Funct. Anal.*, **128**, 163-185 (1995)

[33] Pérez, C., Torres, R.H.: Sharp maximal function estimates for multilinear singular integrals. *Contemp. Math.*, **320**, 323-331 (2003)

[34] Ruiz, A., Vega, L.: Unique comtinuation for Schrödinger operators with potential in Morrey spaces. Publ. Mat. **35**(1),291-298 (1991)

[35] Stein, E.M.: Singular integral and differentiability properties of functions. Princeton University Press, Princeton (1971)

[36] Tang, L.: Weighted estimates for vector-valued commutators of multilinear operators. *Proc. Roy. Soc. Edinburgh Sect. A*, **138**, 897-922 (2008)

[37] Uchiyama, A.: On the compactness of operators of Hankel type. Tôhoku Math. J. **30**, 163-171 (1978)

[38] Wang, D.H., Zhou, J., Chen, W.Y.: Characterization of CMO via compactness of the commutators of bilinear fractional integral operators. arXiv:1612.01116

[39] Wang, S.: The compactness of the commutator of fractional integral operator (in Chinese). Chin. Ann. Math **8**(A), 475-482 (1987)

College of Mathematics and System Sciences, Xinjiang University, Urumqi 830046
Republic of China
E-mail address: Wangdh1990@126.com; zhoujiang@xju.edu.cn; zhidong1960@163.com